



NP Stocker S B1600

MEDICATED

For Further Manufacture of Feed

For increased rate of weight gain in pasture cattle (slaughter, stocker, feeder cattle, and dairy and beef replacement heifers).

ACTIVE DRUG INGREDIENT:

Lasalocid..... 1,600 g/ton

GUARANTEED ANALYSIS:

Calcium (Min).....	18.00%
Calcium (Max).....	21.60%
Phosphorus (Min).....	1.00%
Salt (Min).....	20.00%
Salt (Max).....	24.00%
Magnesium (Min).....	1.00%
Potassium (Min).....	0.50%
Cobalt (Min).....	9 ppm
Copper (Min).....	1,000 ppm
Iodine (Min).....	100 ppm
Manganese (Min).....	800 ppm
Selenium (Min).....	26 ppm
Zinc (Min).....	3,000 ppm
Vitamin A (Min).....	150,000 IU/lb
Vitamin D-3 (Min).....	15,000 IU/lb
Vitamin E (Min).....	150 IU/lb

INGREDIENTS:

Calcium Carbonate, Salt, Processed Grain By-Products, Monocalcium Phosphate, Dicalcium Phosphate, Molasses Products, Magnesium Oxide, Vegetable Oil, Potassium Chloride, Manganous Oxide, Manganese Sulfate, Zinc Oxide, Zinc Sulfate, Copper Sulfate, Sodium Selenite, Ethylenediamine Dihydroiodide, Calcium Iodate, Cobalt Carbonate, Vitamin A Supplement, Vitamin D-3 Supplement, Vitamin E Supplement, Mineral Oil, Red Iron Oxide and Artificial Flavoring.

DIRECTIONS FOR USE:

Thoroughly mix 500 lbs. of **NP Stocker S B1600** with 1,500 lbs. of non-medicated feed. Feed one pound of this medicated mixture to deliver 200 mg lasalocid. Cattle should receive no less than 60 mg nor more than 300 mg lasalocid per head daily in one pound of feed on a continuous basis for the stated drug claims. Four (4) ounces of **NP Stocker S B1600** contains 200 mg lasalocid.

For additional information, please contact your Nutrition Plus LLP representative.

INDICATIONS FOR USE: Intake of lasalocid in excess of 200 mg/head/day have not been shown to be more effective than 200 mg/head/day.

CAUTION: Use as directed. The safety of lasalocid for use in unapproved species has not been established. Do not allow horses or other equines access to feeds containing lasalocid as ingestion may be fatal. Mix thoroughly with grain and/or roughage prior to feeding. Feeding undiluted or mixing errors may result in an excess lasalocid concentration that could be fatal to cattle and sheep. Consumption of selenium should not exceed 3 mg per head daily.

WARNING: A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal. This product, which contains added copper, should not be fed to sheep or any species with a low tolerance to supplemental copper.

NET WEIGHT: 50 LB (22.7 KG)

Manufactured for:
NUTRITION PLUS LLP
 2308 10th Ave.
 Canyon, TX 79015

Batch #

21753