



NP Stocker S Hi Mag Bloat Prevention MEDICATED

For Further Manufacture of Feed

For prevention of legume (alfalfa, clover) and wheat pasture bloat when fed continuously during exposure to bloat-producing conditions.

ACTIVE DRUG INGREDIENT:

Poloxalene 4.4%

GUARANTEED ANALYSIS:

Calcium (Min)	12.0%
Calcium (Max)	14.4%
Phosphorus (Min)	1.0%
Salt (Min)	13.5%
Salt (Max)	16.2%
Magnesium (Min)	6.0%
Cobalt (Min)	9 ppm
Copper (Min)	1,100 ppm
Iodine (Min)	125 ppm
Manganese (Min)	500 ppm
Selenium (Min)	9 ppm
Zinc (Min)	3,000 ppm
Vitamin A (Min)	100,000 IU/lb
Vitamin D-3 (Min)	20,000 IU/lb
Vitamin E (Min)	100 IU/lb

INGREDIENTS:

Calcium Carbonate, Salt, Processed Grain By-Products, Molasses Products, Roughage Products, Monocalcium Phosphate, Dicalcium Phosphate, Magnesium Oxide, Plant Protein Products, Manganous Oxide, Manganese Sulfate, Zinc Oxide, Zinc Sulfate, Copper Sulfate, Sodium Selenite, Ethylenediamine Dihydroiodide, Calcium Iodate, Cobalt Carbonate, Vitamin A Supplement, Vitamin D-3 Supplement, Vitamin E Supplement, Mineral Oil, Red Iron Oxide and Artificial Flavoring.

FEEDING DIRECTIONS:

Feed poloxalene at a rate of 1-2 g per 100 lb. body weight per day for the above drug claim. Eight (8) to sixteen (16) ounces of **NP Stocker S Hi Mag Bloat Prevention** will treat a 1000-lb. animal at the stated rate. For optimum protection against bloat, begin feeding at least 3 days prior to grazing bloat-producing pastures. Provide one covered mineral feeder for each 5-10 animals; and locate mineral feeders where animals congregate (watering and loafing areas).

It is essential that each animal consume the total recommended daily dosage of poloxalene for best bloat protection. Mineral consumption levels will vary depending upon the mineral status of the animals, pasture conditions and animal feeding habits. If the average daily intake of **NP Stocker S Hi Mag Bloat Prevention** drops significantly below expected, be prepared immediately to relocate and increase average daily intake. If animals do not respond immediately with an increase in daily intake, remove the animals from bloat-producing pasture until the proper daily intake is re-established.

Care should be taken to supply a clean, fresh source of water to all animals.

For additional information, please contact your Nutrition Plus LLP representative.

CAUTION: Use as directed. Consumption of selenium should not exceed 3 mg per head daily.

WARNING: This product, which contains added copper, should not be fed to sheep or any species that have a low tolerance to supplemental copper.

NET WEIGHT: 40 LBS. (18.17 KG)

Manufactured for:
NUTRITION PLUS LLP
2308 10th Ave.
Canyon, TX 79015

Batch #



36413